Applicant: Arie Pieter Otte Serial No.: 09/762,916

Filed: June 29, 2001

Page 2

Amendments to the Claims

Please cancel claims 40-42 without disclaimer or prejudice to applicant's right to pursue the subject matter of these claims in a future continuation or divisional application.

1-24. (Canceled)

- 25. (Previously presented) A method of detecting, and optionally selecting, a DNA sequence, wherein the DNA sequence to be detected possesses a stable expression-modulating quality, which method comprises the steps of:
- 1) cloning in a vector of DNA fragments between i) a DNA sequence involved in the induction of gene-transcription repressing chromatin, and ii) a reporter gene comprising a promotor, resulting in a variety of a fragment-comprising vectors;
 - 2) introducing the vectors into a transcription system; and
- 3) subjecting the host cells to a selection step in order to identify the DNA sequence with a stable expression modulating quality;

wherein the DNA sequence involved with the induction of gene-transcription repressing chromatin is a DNA sequence that is recognized by a heterochromatin-binding protein comprising HP1, which HP1-comprising complex is present in the transcription system and/or the host cell.

26. (Previously presented) A method according to claim 25, wherein the DNA sequence comprises an expression-enhancing quality.

Filed: June 29, 2001

Applicant: Arie Pieter Otte

Serial No.: 09/762,916

Page 3

27. (Previously presented) A method according to claim 26, wherein the

transcription system comprises host cells.

28. (Previously presented) A method according to claim 25, wherein the cloned

DNA fragments have a size of 5,000 base pairs.

29. (Previously presented) A method according to claim 25, wherein the distance

between the DNA sequence involved in gene repressing chromatin and the reporter gene

is fewer than 5,000 base pairs.

30. (Previously presented) A method according to claim 25, wherein the

promoter may be active in the transcription system but wherein induction of

gene-repressing chromatin in the vectors results in the repression of the transcription of

the reporter gene.

31. (Previously presented) A method according to claim 25, wherein the selection

in step 3) occurs by using a reporter gene which provides resistance to a growth

inhibitor.

32. (Previously presented) A method according to claim 31, wherein the host

cells are cultivated in the presence of the growth inhibitor.

33. (Previously presented) A method according to claim 32, wherein the growth

inhibitor is present in a concentration sufficiently high to kill host cells in which the gene

providing resistance to the growth inhibitor is not active.

Applicant: Arie Pieter Otte

Serial No.: 09/762,916

Filed: June 29, 2001 Page 4

34. (Previously presented) A method according to claim 33, wherein an antibiotic

is used as the growth inhibitor and the reporter gene provides resistance to the

antibiotic.

35. (Previously presented) A method according to claim 34, wherein the reporter

gene codes for Green Fluorescent Protein.

36. (Previously presented) A method according to claim 35, wherein the reporter

gene is luciferase.

37. (Previously presented) A method according to claim 36, wherein the

fluorescent host cells are separated from non-fluorescent host cells by means of a

Fluorescence-Activated Cell Sorter (FAGS).

38. (Previously presented) A method according to claim 29, wherein the cloned

DNA fragments have a size of substantially between 2,000-3,000 base pairs.

39-42. (Canceled)

43. (Previously presented) A method according to claim 25, wherein the DNA

sequence involved with the transcription inducing of gene-repressing chromatin is a DNA

sequence that is selectively recognized by at least one DNA-binding protein and the

organism also expresses a protein complex comprising i) a first part selectively binding

the DNA sequence, and ii) a second part inducing the formation of chromatin in which

the transcription is repressed.

Applicant: Arie Pieter Otte

Serial No.: 09/762,916

Filed: June 29, 2001 Page 5

44. (Previously presented) A method according to claim 43, wherein the protein

complex comprises a fusion protein.

45. (Previously presented) A method according to claim 44, wherein first part is a

part binding the DNA binding site of LexA-DNA or GAL4-DNA.

46. (Previously presented) A method according to claim 25, wherein the

organism in step 1) is selected from the group comprising a plant and a vertebrate.

47. (Previously presented) A method according to claim 46, wherein the

vertebrate is a mammal.

48. (Previously presented) A method according to claim 25, wherein the vector is

an episomally replicating vector.

49. (Previously presented) A method according to claim 48, wherein the vector

comprises a replication origin from the Epstein-Barr virus (EBV), OriP, and a nuclear

antigen (EBNA1).

50-55. (Canceled)